

# Subriemannian Geometry: The basic notations and examples

Winterschool in Geilo, Norway

Wolfram Bauer

Leibniz U. Hannover

March 4-10. 2018

## Outline

1. Motivations, definitions and examples
2. Horizontal curves and an optimal control problem
3. More examples and constructions

## Motivation: Sub-Riemannian geometry

Consider  $n$  classical particles with **coordinates**  $\{q_1, \dots, q_n\}$ .

### Motion under constraints

H:  $f(q_1, \dots, q_n) = 0$ , (*holonomic*),

NH:  $f(q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_n) = 0$ , (*non-holonomic*).

### Exampels:

H: A particle moving along a **surface**, or a **pendulum**.

NH: Rolling of a ball on a plane (or some surface) **without slipping or twisting**.

### Corresponding geometric structures on a manifold

- *holonomic constraints*  $\longrightarrow$  integrable distribution (foliation of a manifold),
- *non-holonomic constraints*  $\longrightarrow$  **Sub-Riemannian structure**.

## Parking a car: Rototranslation

Position of the car robot in **3-space**:  $(x, y, \vartheta) \in \mathbb{R}^2 \times \mathbb{S}^1$ .

### Possible movements

- $X = \cos \vartheta \cdot \partial_x + \sin \vartheta \cdot \partial_y$ , (in direction of the car)
- $Y = \partial_\vartheta$ , (rotation)
- $Z = -\sin \vartheta \cdot \partial_x + \cos \vartheta \cdot \partial_y$ , (orthogonal to the car).

## Parkin a car: Rototranslation

**Connecting positions:** Which movements allow to reach from any position of the car any other position?

### Observations

- Moving only along  $X$  and  $Z$  is **not enough**: it keeps the angle  $\vartheta$  fixed.

$$\text{span}\{X, Z\} = \ker d\vartheta \quad \text{and} \quad d\vartheta = \text{closed form}, \\ [X, Z] = 0.$$

- Moving along  $X$  and  $Y$  (*parking procedure*) **might be sufficient** for connecting positions.

$$\text{span}\{X, Y\} = \ker \omega \quad \text{where} \quad \omega = -\sin \vartheta dx + \cos \vartheta dy. \\ [X, Y] = \left[ \cos \vartheta \cdot \partial_x + \sin \vartheta \cdot \partial_y, \partial_\vartheta \right] \\ = -\sin \vartheta \cdot \partial_x + \cos \vartheta \cdot \partial_y = Z.$$

## Sub-Riemannian Geometry

*"Sub-Riemannian geometry models motions under non-holonomic constraints".*

### Definition

A **Sub-Riemannian manifold** (shortly: SR-m) is a triple  $(M, \mathcal{H}, \langle \cdot, \cdot \rangle)$  with:

- $M$  is a smooth manifold (without boundary),  $\dim M \geq 3$  and  $\mathcal{H} \subset TM$  is a **vector distribution**.
- $\mathcal{H}$  is **bracket generating** of rank  $k < \dim M$ , i.e.

$$\text{Lie}_x \mathcal{H} = T_x M.$$

- $\langle \cdot, \cdot \rangle_x$  is a smoothly varying family of inner products on  $\mathcal{H}_x$  for  $x \in M$ .

# 1.Example: Heisenberg group

Consider the 3- dimensional **Heisenberg group**  $\mathbb{H}_3 \cong (\mathbb{R}^3, *)$  with **product**:

$$(x_1, y_1, z_1) * (x_2, y_2, z_2) = \left( x_1 + x_2, y_1 + y_2, z_1 + z_2 + \frac{1}{2}[x_1 y_2 - y_1 x_2] \right).$$

**Lie algebra of  $\mathbb{H}_3$ :**

On  $\mathbb{H}_3 \cong \mathbb{R}^3$  define **left-invariant vector fields**: Let  $q = (x, y, z) \in \mathbb{H}_3$ : <sup>1</sup>

$$\begin{aligned} [X_1 f](q) &= \frac{df}{dt} \left( q * (t, 0, 0) \right) \Big|_{t=0} \\ &= \frac{df}{dt} \left( x + t, 0, z - \frac{yt}{2} \right) = \left[ \left( \frac{\partial}{\partial x} - \frac{y}{2} \frac{\partial}{\partial z} \right) f \right] (q). \end{aligned}$$

Similarly, with curves  $(0, t, 0)_t$  and  $(0, 0, t)_t$ :

$$X_2 = \frac{\partial}{\partial y} + \frac{x}{2} \frac{\partial}{\partial z} \quad \text{and} \quad Z = \frac{\partial}{\partial z}.$$

<sup>1</sup>"X left-invariant":  $X_{g*h} = (L_g)_* X_h$  with the left-multiplication  $L_g : \mathbb{H}_3 \rightarrow \mathbb{H}_3$ .

## Heisenberg group as SR-manifold

**Known fact:**

The **Lie algebra**  $(\mathfrak{h}_3, [\cdot, \cdot])$  of  $\mathbb{H}_3$  can be identified with:

$$\mathfrak{h}_3 = \text{span} \{ X_1, X_2, Z \} \quad \text{with} \quad [\cdot, \cdot] = \text{commutator of vector fields.}$$

### Observation

If we calculate **Lie-brackets**  $[\cdot, \cdot]$ , then one only finds one non-trivial bracket relation is:

$$[X_1, X_2] = X_1 X_2 - X_2 X_1 = Z.$$

- Put  $\mathcal{H} = \text{span} \{ X_1, X_2 \} \subset T\mathbb{H}_3$  (distribution),
- Define  $\langle \cdot, \cdot \rangle$  on  $\mathcal{H}$  by **declaring**  $X_1$  and  $X_2$  pointwise orthonormal.

**Conclusion:**  $(\mathbb{H}_3, \mathcal{H}, \langle \cdot, \cdot \rangle)$  defines a Sub-Riemannian structure on  $\mathbb{H}_3$ .

## Horizontal curves and cc-distance:

On a SR-manifold  $(M, \mathcal{H}, \langle \cdot, \cdot \rangle)$  we consider **horizontal objects**, i.e. objects under non-holonomic constraints.

### Example

Consider a curve  $\gamma : [0, 1] \rightarrow M$ :<sup>a</sup>

- $\gamma$  is called **horizontal**, (a.e.) it is **tangential** to  $\mathcal{H}$ , i.e.

$$\dot{\gamma}(t) \in \mathcal{H}_{\gamma(t)}.$$

- The **curve length** is defined by:

$$\ell(\gamma) := \int_0^1 \sqrt{\langle \dot{\gamma}(t), \dot{\gamma}(t) \rangle_{\gamma(t)}} dt.$$

- **SR geodesic** = locally length minimizing horizontale curve.

---

<sup>a</sup>piecewise  $C^1$  or just absolutely continuous

## Carnot-Carathéodory metric

**Definition:** Sub-Riemannian distanced (cc-distance)

The **SR distance** between two points  $a, b \in M$  is defined by:

$$d_{cc}(a, b) := \inf \left\{ \ell(\gamma) : \gamma \text{ horizontal}, \gamma(0) = a, \gamma(1) = b \right\}.$$

**Question:** Let  $M$  be a connected SR-manifold. Can we connect any two points on  $M$  by **horizontal curves**?

**Theorem** (W.-L. Chow 1939, P.-K. Rashevskii 1938)

Any two points on a connected SR-manifold can be connected by **piecewise smooth horizontal curves**.

**Consequence:** The cc-distance  $d_{cc}$ <sup>2</sup> on a connected SR-manifold is **finite**. Hence  $(M, d_{cc})$  forms a **metric space**.

---

<sup>2</sup>Carnot-Carathéodory distance

# Geodesic equations

## Some question:

- How can we **obtain** Sub-Riemannian geodesics?
- Relation to  $d_{cc}$ : can we realize the cc-distance between two point by a **(piecewise) smooth SR geodesic**?
- Is the distance  $x \mapsto d_{cc}(x_0, x)$  **smooth** for fixed points  $x_0$ ?

Let  $(M, \mathcal{H}, \langle \cdot, \cdot \rangle)$  be a SR-manifold. Let

$$[X_1, \dots, X_m] = \text{vector fields} \quad \text{and} \quad m = \text{rank } \mathcal{H}.$$

an **local orthonormal frame** around a point  $q \in M$ , i.e.

$$\mathcal{H}_q = \text{span} \{ X_1(q), \dots, X_m(q) \} \quad \text{and} \quad \langle X_i(q), X_j(q) \rangle = \delta_{ij}.$$

**Idea:** *Expand locally the derivative of a horizontal curve with respect to the above frame*

## SR-geodesics and optimal control

### Observation

Let  $\gamma : [0, 1] \rightarrow M$  be **horizontal**. With suitable **coefficients**  $u_j(t)$  one can write

$$\gamma'(t) = \sum_{j=1}^m u_j(t) \cdot X_j(t) \quad \implies \quad \langle \gamma'(t), \gamma'(t) \rangle = \sum_{j=1}^m u_j^2(t).$$

Finding SR-geodesics between  $A, B \in M =$  **optimal control problem OCP**.

**OCP:** *Minimize the cost*

$$J_T(u) := \frac{1}{2} \int_0^T \sqrt{\sum_{j=1}^m u_j^2(t)} dt$$

*under the conditions*

$$\gamma' = \sum_{j=1}^m u_j \cdot X_j(\gamma) \quad \text{and} \quad \gamma(0) = A, \gamma(T) = B.$$

## SR-geodesic: a Hamiltonian formalism

### Remark:

Instead of minimizing a length we may equivalently minimize an "energy":

**OCP:** Minimize the cost

$$J_T(u) := \frac{1}{2} \int_0^T \sum_{j=1}^m u_j^2(t) dt$$

under the conditions

$$\gamma' = \sum_{j=1}^m u_j \cdot X_j(\gamma) \quad \text{and} \quad \gamma(0) = A, \gamma(T) = B.$$

**Hamiltonian formalism** (as known in Riemannian geometry):

Assign a **Sub-Riemannian Hamiltonian**  $H_{sr} \in C^\infty(T^*M)$  to the problem:

$$H_{sr}(q, p) = \sum_{j=1}^m p(X_j(q))^2 \quad \text{where} \quad (q, p) \in T_q^*M.$$

## SR-geodesic: a Hamiltonian formalism

With the Poisson bracket  $\{\cdot, \cdot\}$  on  $C^\infty(T^*M)$  consider:

$$\vec{H}_{sr} = \{\cdot, H\} = \frac{\partial H}{\partial p} \cdot \frac{\partial}{\partial q} - \frac{\partial H}{\partial q} \cdot \frac{\partial}{\partial p} = \text{Hamiltonian vector field}$$

The Hamiltonian vector field defines the **geodesic flow** on  $T^*M$  and projections of the flow to  $M$  give SR-geodesics:

### Theorem (normal geodesics)

Let  $\zeta(t) = (\gamma(t), p(t))$  be a solution to the **normal geodesic equations**:

$$\dot{q} = \frac{\partial H}{\partial p_i}(q, p) \quad \text{and} \quad \dot{p} = -\frac{\partial H}{\partial q_i}(q, p), \quad i = 1 \dots \dim M.$$

Then  $\gamma(t)$  **locally minimizes** the SR-distance.

### Proof: <sup>3</sup>

<sup>3</sup>R. Montgomery, *A tour of Subriemannian Geometries, Their Geodesics and Applications* Math. Surveys and Monographs, 2002.

# SR-geodesics

## Remark

There are various differences to the setting of a Riemannian manifold:

- The **Hamiltonian in Riemannian geometry** can be expressed as

$$H_r(q, p) = \sum_{i,j=1}^n g^{ij}(q) p_i p_j, \quad g^{ij} := \text{inverse metric tensor.}$$


In SR-geometry  $g_{ij}$  is an  $m \times m$ -matrix and not invertible on  $TM$ .

- There are no **2nd order geodesic equations** in the SR-setting such as

$$\ddot{q}^k = \Gamma_{ij}^k \dot{q}_i \dot{q}_j.$$

The obtained **regularity** of SR-geodesics is not clear.

- In SR-geometry there may be **singular geodesics** which **do not solve** the **geodesic equations** in the above theorem.



**The falling cat:**

A connectivity  
problem  
in SR geometry



# Generalizations of the Heisenberg group

A Lie group  $G$  has **trivial tangent bundle** and the last construction of a trivial bundle can be generalized:

## Left-invariant structure

- Let  $\mathfrak{g}$  denote the Lie algebra of  $G$ .
- Let  $V \subset \mathfrak{g}$  be a subspace of  $\mathfrak{g}$  with **inner product**  $\langle \cdot, \cdot \rangle_V$  and

$$\mathfrak{g} = \text{Lie}(V) = \text{span} \left\{ v, [w, x], [y, [w, x]], \dots : x, y, w \in V \right\}.$$

Identify  $V$  (via left-translation) with a space of **left-invariant vector fields** on  $G$ .

- The  $G$  becomes a sub-Riemannian manifold  $(G, \mathcal{H}, \langle \cdot, \cdot \rangle)$  with:

$$\begin{aligned} \mathcal{H} &= V \\ \langle \cdot, \cdot \rangle_q &= \langle (dL_q)^{-1} \cdot, (dL_q)^{-1} \cdot \rangle_V. \end{aligned}$$

## Contact structures

Let  $\Theta$  be a **one-form** on a manifold  $M$  of dimension  $\dim M = 2k + 1$ . Put:

$$\mathcal{H}_q := \text{kern}(\Theta_q) \subset T_q M, \quad (q \in M).$$

## Contact form

Assume that  $\Theta$  has the following properties:

- the **restriction** of  $d\Theta_q$  to  $\mathcal{H}_q$  is **non-degenerate**<sup>a</sup> for each  $q \in M$ :

*If  $v \in \mathcal{H}$  with  $d\Theta(v, w) = 0$  for all  $w \in \mathcal{H}_q$ , then  $v = 0$ .*

- **equivalently:** the form

$$\omega := \Theta \wedge (d\Theta)^{2k} \neq 0$$

does not vanish at any point of  $M$  ( $= \omega$  is a **volume form**):

---

<sup>a</sup>a symplectic form

# Contact manifolds

## Lemma

Let  $\Theta$  be a **contact form** on  $M$ . Then

$$\mathcal{H} := \ker \Theta \subset TM$$

is a **bracket generating** distribution.

**Proof:** Use **Cartan's formula**:

$$d\Theta(X, Y) = X\Theta(Y) - Y\Theta(X) - \Theta([X, Y]).$$

Let  $X, Y$  be **horizontal**, i.e.  $X_q, Y_q \in \mathcal{H}_q = \ker \Theta_q$  for all  $q \in M$ . Then

$$\Theta(X) = \Theta(Y) = 0 \implies d\Theta(X, Y) = -\Theta([X, Y]).$$

Since  $d\Theta$  is **non-degenerate** on  $\mathcal{H}_q$  we find  $X, Y$  with

$$[X, Y]_q \notin \ker \Theta_q = \mathcal{H}_q.$$

□

## Contact manifolds (continued)

Choose an **almost complex structure**  $J : \mathcal{H} \rightarrow \mathcal{H}$  such that

$$\langle \cdot, \cdot \rangle = d\Theta(J\cdot, \cdot), \quad \text{and} \quad J^2 = -I$$

is an **inner product** on  $\mathcal{H}$  (symmetric, positive definite).

### Definition (contact Sub-Riemannian manifold)

The triple  $(M, \mathcal{H}, \langle \cdot, \cdot \rangle)$  is called **contact Sub-Riemannian manifold**.

**Example:** Consider again the **Heisenberg group**  $\mathbb{H}_3 \cong \mathbb{R}^3$  with distribution:

$$\mathcal{H} = \text{span} \left\{ \frac{\partial}{\partial x} - \frac{y}{2} \frac{\partial}{\partial z}, \frac{\partial}{\partial y} + \frac{x}{2} \frac{\partial}{\partial z} \right\} = \ker \underbrace{\left( dz - \frac{x}{2} dy + \frac{y}{2} dx \right)}_{=\Theta}.$$

Moreover,  $\Theta$  is a contact form and  $\mathbb{H}_3$  is a **contact SR-manifold**:

$$\Theta \wedge d\Theta = -\Theta \wedge (dx \wedge dy) = -dx \wedge dy \wedge dz \neq 0.$$


# Rototranslation group: How to park a car?

## Possible movements

- $X = \cos \vartheta \cdot \partial_x + \sin \vartheta \cdot \partial_y$ , (in direction of the car)
- $Y = \partial_\vartheta$ , (rotation)
- $Z = -\sin \vartheta \cdot \partial_x + \cos \vartheta \cdot \partial_y$ , (orthogonal to the car).

Good choice:

$$\mathcal{H} = \text{span}\{X, Y\} = \text{kern } \omega \quad \text{with} \quad \omega = -\sin \vartheta \cdot dx + \cos \vartheta \cdot dy.$$



$$\omega \wedge d\omega = \omega \wedge (-\cos \vartheta \cdot d\vartheta \wedge dx - \sin \vartheta \cdot d\vartheta \wedge dy) = -dx \wedge dy \wedge d\vartheta \neq 0.$$

## Sub-Riemannian structures on spheres

Different from a Lie group it is well-known that most of the Euclidean unit spheres  $\mathbb{S}^n \subset \mathbb{R}^{n+1}$  of dimension  $n$  do not have a trivial tangent bundle.

### Exceptions

Precisely the spheres  $\mathbb{S}^n$  where  $n = 1, 3, 7$  have trivial tangent bundle.

**Questions:** Are there:

- (1) bracket generating distributions on Euclidean spheres?
- (2) trivializable bracket generating distributions  $\mathcal{H}$  on  $\mathbb{S}^n$ , (i.e.  $\mathcal{H}$  is trivial as a vector bundle)?

**Answers:**

- (1) There are various constructions:
  - ▶ odd dimensional spheres  $\mathbb{S}^{2k+1} \subset \mathbb{R}^{2n} \cong \mathbb{C}^n$  carry a contact structure (from the diagonal action of  $S^1$  on  $\mathbb{C}^n$ ),
  - ▶ via (quaternionic) Hopf fibration in some dimensions, ...
- (2) In some dimensions via canonical vector fields.

## SR-structures on spheres

There are various constructions of SR-structures on **Euclidean spheres**. Some models arise from different points of view, e.g.  $\mathbb{S}^3$  or  $\mathbb{H}_3$  are:

*Lie groups, total space of a fiber bundle (e.g. Hopf fibration), contact manifolds, ...*



W. -B. K. Furutani, C. Iwasaki

*Trivializable sub-Riemannian structures on spheres*, Bull. Sci. math. 137 (2013), 361- 385.



O. Calin, D.-C. Chang,

*Sub-Riemannian geometry on the sphere  $S^3$* , Canad. J. Math. 61 (4) (2009) 721 - 739.



I. Markina, M.G. Molina,

*Sub-Riemannian geodesics and heat operator on odd dimensional spheres*, Anal. Math. Phys. 2 (2) (2012) 123 - 147

## Adams Theorem

Theorem (J.F. Adams, 1962)

The **maximal dimension**  $\gamma(n)$  of a **trivial subbundle** in  $T\mathbb{S}^n$  is:

$$\gamma(n) = 2^a + 8b - 1.$$

The numbers  $0 \leq a < 4$  and  $0 \leq b$  are determined through the relations:

$$n + 1 = 2^{a+4b} \times [\text{odd}].$$

**Canonical vector fields:** For  $\alpha = 1, \dots, \gamma(n)$  consider:

$$X_\alpha := \sum_{i=1}^{n+1} \sum_{j=1}^{n+1} a_{ij}^\alpha x_i \frac{\partial}{\partial x_j}, \quad \text{with} \quad A_\alpha = (a_{ij}^\alpha) \in \mathbb{R}(n+1).$$

Assume that the matrices  $A_\alpha$  fulfill the **Clifford relations**:

$$A_\alpha A_\beta + A_\beta A_\alpha = -2\delta_{\alpha\beta} I.$$

# Sub-Riemannian structures via canonical vector fields

## Lemma

The restriction of the **canonical vector fields** to  $\mathbb{S}^n$  are orthonormal at each point of  $\mathbb{S}^n$ . The distribution:

$$\mathcal{H} = \text{span}\left\{X_\alpha : \alpha = 1, \dots, \gamma(n)\right\}$$

defines a **maximal dimensional trivial subbundle** of  $T\mathbb{S}^n$ .

The **Clifford relations** imply relations on the **brackets** of canonical vector fields. In particular, these show:

$$[X_\alpha [X_\beta [X_\gamma \dots]]] \in \text{span}\left\{X_i, [X_j, X_k] : i, j, k = 1, \dots, \gamma(n)\right\}.$$

**Necessary condition for the bracket generating property:**

$$\rho(n) := \gamma(n) + \binom{\gamma(n)}{2} > n. \quad (*)$$

## Trivializable Sub-Riemannian structures on sphere

**Lemma** Property (\*) precisely holds in the following dimensions:

$n$	1	3	7	15	23	31	63
$\gamma(n)$	1	3	7	8	7	9	11
$\rho(n)$	1	6	28	36	28	45	66

**Next task:** Sufficient conditions for the bracket generating property.

Theorem, (B. Furutani, Iwasaki)

**Trivializable Sub-Riemannian structures** on spheres  $\mathbb{S}^n$  via a Clifford module structure **only exist** in the following dimensions:


$$n = 3, 7, 15.$$

On  $\mathbb{S}^7$  there are trivializable structures of rank 4, 5, 6.

**Question:** Are there Subriemannian structures on **exotic 7-spheres**?

# Gromoll-Meyer exotic 7-sphere $\Sigma_{GM}^7$

Exotic 7-sphere as base of a  $\Delta$ -principal bundle



With  $\Delta = \{(\lambda, \lambda) : \lambda \in \text{Sp}(1)\}$  und  $G = \text{Sp}(1) \times \text{Sp}(1) \supset \Delta$ .

Theorem (B., Furutani, Iwasaki, 2016)

The bi-quotient of compact groups induces a **rank 4 SR-structure** on the Gromoll-Meyer exotic 7-sphere.

$$T(\text{Sp}(2)) = V^\Delta \oplus H^\Delta = V^G \oplus H^G \text{ und } H^G \subset H^\Delta.$$

## Sub-Riemannian structures of bundle type

Let  $(M, g_M)$  and  $(N, g_N)$  be **Riemannian manifolds** with **Riemannian submersion**:

$$\pi : M \rightarrow N.$$

### Properties

Let  $q \in M$  and  $p = \pi(q) \in N$ .

- kern  $d\pi_q \subset T_q M$  is a the space tangent to the fibre  $\pi^{-1}(p)$  at  $q$ .
- The **restriction** of the differential

$$d\pi_q : \mathcal{H}_q := (\text{kern } d\pi_q)^\perp \subset T_q M \rightarrow T_p N$$

is an **isometry**.

- On  $\mathcal{H}$  consider the restriction  $\langle \cdot, \cdot \rangle$  of the metric on  $TM$

These data may give a **SR-structure of bundle type**. (Note: bracket generating property is not clear in general and has to be checked).

## Example: Hopf fibration

Consider the three sphere as a subset of  $\mathbb{C}^2$ :

$$\mathbb{S}^3 = \{z = (z_1, z_2) \in \mathbb{C}^2 : |z_1|^2 + |z_2|^2 = 1\} \subset \mathbb{C}^2.$$

### Definition (Hopf fibration)

The **Hopf fibration** is the submersion map

$$\pi : \mathbb{S}^3 \rightarrow \mathbb{S}^2_{\frac{1}{2}} : \pi(z) := \frac{1}{2} \left( |z_1|^2 - |z_2|^2, \operatorname{Re}(z_1 \bar{z}_2), \operatorname{Im}(z_1 \bar{z}_2) \right),$$

where  $\mathbb{S}^2_{\frac{1}{2}}$  is the 2-sphere of radius  $1/2$ .






**Theorem:** The **Hopf fibration** defines a **principal  $\mathbb{S}^1$ -bundle**, where  $\mathbb{S}^1$  act by componentwise multiplication on  $\mathbb{S}^3 \subset \mathbb{C}^2$ .

**Remark:** The corresponding distribution on  $\mathbb{S}^3$  of bundle type is bracket generating (and coincides with a **contact structure** on  $\mathbb{S}^3$ ).




## Summary

- Sub-Riemannian geometry models motion under non-holonomic constraints (mechanical systems, rolling of manifolds, parking a car, falling cat...)
- Connected SR-manifolds are metric spaces with the cc-distance.
- Sub-Riemannian geodesics  $\longleftrightarrow$  optimal control problem.
- Examples include: some Lie groups, (e.g. Heisenberg group or  $\mathbb{S}^3$ ), Euclidean spheres, some principal bundles (e.g. Hopf fibration).

## References

-  [W. Bauer, K. Furutani, C. Iwasaki,](#)  
*A codimension 3 sub-Riemannian structure on the Gromoll-Meyer exotic sphere*, Differential geometry and its applications, 53 (2017), 114-136.
-  [W. -B. K. Furutani, C. Iwasaki](#)  
*Trivializable sub-Riemannian structures on spheres*, Bull. Sci. math. 137 (2013), 361- 385.
-  [O. Calin, D.-C. Chang](#)  
*Sub-Riemannian Geometry - General Theory and Examples*, Cambridge University Press, 2009.
-  [A. Bellaïche](#)  
*Sub-Riemannian geometry*, Basel-Boston-Berlin, 1996.
-  [M. Gromov,](#)  
*Carnot-Carathéodory spaces seen from within*, Sub-Riemannian Geometry, Birkhäuser, Progress in Math. 144, (1996), 79-323.

## References

-  [I. Markina, M.G. Molina,](#)  
*Sub-Riemannian geodesics and heat operator on odd dimensional spheres*, Anal. Math. Phys. 2 (2) (2012) 123 - 147
-  [I. Markina,](#)  
*Geodesics in geometry with constraints and applications. Quantization, PDEs, and geometry*, 153- 314, in: Oper. Theory Adv. Appl., 251, Adv. Partial Differ. Equ. (Basel), Birkhäuser/Springer, Cham, 2016.
-  [R. Montgomery,](#)  
*A tour of Subriemannian Geometries, Their Geodesics and Applications*, Mathematical Surveys and Monographs, 91, 2002.



**Thank you for your attention!**