

# Lecture I: Introduction to Special Geometry

Vicente Cortés  
Department of Mathematics  
University of Hamburg

Winter School “Geometry, Analysis, Physics”  
Geilo (Norway), March 4-10, 2018

# Plan of the mini course

- I. Introduction to special geometry
- II. Geometric constructions relating different special geometries I
- III. Geometric constructions relating different special geometries II
- IV. Completeness results
- V. Construction of complete quaternionic Kähler manifolds

## Plan of the first lecture

- ▶ Physical motivation of special geometry
- ▶ Affine and projective special real geometry
- ▶ Affine and projective special Kähler geometry
- ▶ Hyper-Kähler and quaternionic Kähler geometry

## Some references for Lecture I

[CNS] C.–, Nardmann, Suhr (CAG '16)

[AC] Alekseevsky, C.– (CMP '09).

[CM] C.–, Mohaupt (JHEP '09).

[CMMS] C.–, Mayer, Mohaupt, Saueressig (JHEP '04).

[ACD] Alekseevsky, C.–, Devchand (JGP '02).

[F] Freed (CMP '99).

[L] LeBrun (Duke '91).

[GST] Günaydin, Sierra, Townsend (NPB '84).

[DV] de Wit, Van Proeyen (NPB '84).

[BW] Bagger, Witten (NPB '83)

# Physical motivation

## Scalar geometry

$$\mathcal{L} = - \sum g_{ij}(\phi^1, \dots, \phi^n) h^{\mu\nu} \partial_\mu \phi^i \partial_\nu \phi^j + \dots$$

## Physics Definition

**Special geometry** is the scalar geometry of supersymmetric field theories with 8 real supercharges.

## One distinguishes between

- ▶ **Affine** special geometry/supersymmetric **gauge theories** and
- ▶ **Projective** special geometry/**supergravity** theories.

# Affine special real manifolds I: extrinsic and intrinsic definition

## Definition

An **affine special real (ASR) manifold** is a domain  $M \subset \mathbb{R}^n$  endowed with a Riemannian metric  $g = \partial^2 h$ , which is the Hessian of a cubic polynomial.

## Definition

An **intrinsic ASR manifold**  $(M, \nabla, g)$  is a Riemannian manifold  $(M, g)$  endowed with a flat torsion-free connection  $\nabla$  such that  $\nabla g$  is completely symmetric and  $\nabla$ -parallel.

## Remark

ASR mfs. = scalar mfs. of 5d vector multiplets [CMMS]

# Affine special real manifolds II: intrinsic characterization

## Theorem [AC]

- (i) Let  $(M, g)$  be an  $n$ -dim. **ASR manifold** with flat connection  $\nabla$  induced from the inclusion  $M \subset \mathbb{R}^n$ . Then  $(M, \nabla, g)$  is an **intrinsic ASR manifold**.
- (ii) Conversely, let  $(M, g, \nabla)$  be an  $n$ -dim. s.c. **intrinsic ASR manifold**. Then there exists an affine immersion  $\varphi : (M, \nabla) \rightarrow \mathbb{R}^n$  and a cubic polynomial  $h$  s.t.  $g = \varphi^* \partial^2 h$ . In particular,  $(\varphi(M), \partial^2 h)$  is an **ASR manifold**. ( $\varphi$  is unique up to affine transformations of  $\mathbb{R}^n$ .)

# Projective special real manifolds I: extrinsic definition

## Definition

A **projective special real (PSR) manifold** is a hypersurface  $\mathcal{H} \subset \mathbb{R}^{n+1}$  s.t.  $\exists$  homog. cubic polynomial  $h$  on  $\mathbb{R}^{n+1}$  s.t.

- i)  $h = 1$  on  $\mathcal{H}$  and
- ii)  $\partial^2 h$  is negative definite on  $T\mathcal{H}$ .

$\mathcal{H}$  is endowed with the Riemannian metric

$$g_{\mathcal{H}} = -\frac{1}{3} \iota^* \partial^2 h,$$

where  $\iota : \mathcal{H} \rightarrow \mathbb{R}^{n+1}$  is the inclusion map.

$\mathcal{H}$  **complete** :  $\iff (\mathcal{H}, g_{\mathcal{H}})$  complete.

## Remark

PSR mfs. = scalar mfs. of 5d sugra coupled to vector multpl. [GST]

# Centroaffine structures

## Definition

A **centroaffine structure** on a smooth manifold  $M$  is a triple  $(\nabla, g, \nu)$  consisting of a torsion-free connection  $\nabla$ , a pseudo-Riemannian metric  $g$  and a volume form  $\nu$  such that

- (i)  $\nabla\nu = 0$ ,
- (ii) the curvature  $R$  of  $\nabla$  is given by

$$R(X, Y)Z = -(g(Y, Z)X - g(X, Z)Y)$$

for all  $X, Y, Z \in \mathfrak{X}(M)$  and

- (iii)  $\nabla g$  is completely symmetric.

$(M, \nabla, g, \nu)$  is called a **centroaffine manifold**.



# Induced centroaffine structure on a hypersurface

Let  $\mathcal{H} \subset (\mathbb{R}^{n+1}, \det)$  be a **centroaffine hypersurface**, i.e. a nondeg. hypersurface transversal to the position vector field  $\xi$ .

Then

- (i)  $\nu = \iota_\xi \det$  is a volume form on  $\mathcal{H}$  and
- (ii) the (affine) Gauß equation

$$\partial_X Y = \nabla_X Y + g(X, Y)\xi, \quad X, Y \in \mathfrak{X}(\mathcal{H}),$$

defines a centroaffine structure  $(\nabla, g, \nu)$ .

## Remarks

1. The metric  $g$  is called the **centroaffine metric**.
2. For a PSR manifold  $(\mathcal{H}, g_{\mathcal{H}})$  we have  $g = g_{\mathcal{H}}$ .

## Projective special real manifolds II: intrinsic definition

### Definition

An **intrinsic PSR manifold** is a centroaffine manifold  $(M, \nabla, g, \nu)$  with  $g > 0$  such that the covariant derivative of the cubic form  $C = \nabla g$  is given by

$$\begin{aligned}(\nabla_X C)(Y, Z, W) = \\ g(X, Y)g(Z, W) + g(X, Z)g(W, Y) + g(X, W)g(Y, Z),\end{aligned}$$

for all  $X, Y, Z, W \in \mathfrak{X}(M)$ .

### Remark

The above equation implies that  $\nabla C$  is totally symmetric, that is a quartic form.

# Projective special real manifolds III: intrinsic characterization

## Theorem [CNS]

- (i) Let  $\mathcal{H} \subset \mathbb{R}^{n+1}$  be a **PSR manifold** with its induced centroaffine structure  $(\nabla, g, \nu)$ . Then  $(\mathcal{H}, \nabla, g, \nu)$  is an **intrinsic PSR manifold**.
- (ii) Conversely, let  $(M, \nabla, g, \nu)$  be a s.c. **intrinsic PSR manifold**. Then there exists an immersion  $\varphi : M \rightarrow \mathbb{R}^{n+1}$  such that  $\mathcal{H} := \varphi(M) \subset \mathbb{R}^{n+1}$  is a **PSR manifold**.

The immersion  $\varphi$  is unique up to linear unimodular transformations of  $\mathbb{R}^{n+1}$ .

# Affine special Kähler manifolds

## Definition

A (pseudo-) Kähler manifold  $(M, g, J)$  is a (pseudo-) Riemannian manifold  $(M, g)$  endowed with a parallel skew-symm. cx. str.  $J$ .

## Definition [F]

An affine special (pseudo-) Kähler manifold  $(M, J, g, \nabla)$  is a (pseudo-) Kähler mf.  $(M, J, g, \nabla)$  endowed with a flat torsionfree connection  $\nabla$  such that

- (i)  $\nabla\omega = 0$ , where  $\omega = g(\cdot, J\cdot)$ ,
- (ii)  $d^\nabla J = 0$ , where  $J$  is considered as a 1-form with values in  $TM$ .

## Remark

Affine special Kähler mfs. = scalar mfs. of 4d N=2 vector multiplets,

Projective special Kähler mfs. = scalar mfs. of 4d supergravity coupled to N=2 vector multiplets [DV].

# Conical and projective special Kähler manifolds

## Definition [ACD, CM]

A **conical affine special Kähler (CASK) manifold**  $(M, J, g, \nabla, \xi)$  is an affine special (pseudo-)Kähler manifold  $(M, J, g, \nabla)$  endowed with a vector field  $\xi$  such that

- (i)  $\nabla\xi = D\xi = \text{Id}$ , where  $D$  is the Levi Civita connection and
- (ii)  $g$  is positive definite on  $\mathcal{D} := \text{span}\{\xi, J\xi\}$  and negative definite on  $\mathcal{D}^\perp$ .

$\Rightarrow \xi$  and  $J\xi$  generate a hol. action of a 2-dim. Abelian Lie algebra. We will assume that the action lifts to a principal  $\mathbb{C}^*$ -action with the base  $\bar{M} = M/\mathbb{C}^*$ . Then  $J\xi$  generates a free isometric and Hamiltonian  $S^1$ -action and  $\bar{M}$  inherits a Kähler metric  $\bar{g}$ .  $(\bar{M}, \bar{g})$  is called a **projective special Kähler (PSK) manifold**.

# Extrinsic construction of special Kähler manifolds I

## The ambient space

$V = (\mathbb{C}^{2n}, \Omega, \tau)$ ,  $\Omega = \sum dz^i \wedge dw_i$ ,  $\tau = \text{cx. conjugation}$ .

$\rightarrow$  pseudo-Hermitian form  $\gamma := \sqrt{-1}\Omega(\cdot, \tau\cdot)$ .

## Definition

A holomorphic immersion  $\phi : M \rightarrow V$  is called **nondegenerate** if  $\phi^*\gamma$  is nondeg. It is called **Lagrangian** if  $\phi^*\Omega = 0$  and  $\dim M = n$ .

## Theorem [ACD]

- ▶ A nondeg. hol. Lagrangian immersion  $\phi : M \rightarrow V$  induces an affine special pseudo-Kähler structure  $(J, g, \nabla)$  on  $M$ .
- ▶ Every s.c. affine special (pseudo-) Kähler mf.  $(M, J, g, \nabla)$  of dim.  $n$  admits a nondeg. Lagr. immersion  $\phi : M \rightarrow V$  inducing  $(J, g, \nabla)$  on  $M$ . The immersion is unique up to affine transformations with real symplectic linear part.

## Extrinsic construction of special Kähler manifolds II

### Example (affine special pseudo-Kähler domains)

Let  $F$  be a holomorphic function defined on a domain  $M \subset \mathbb{C}^n$  such that the matrix

$$(N_{ij}) = (2\text{Im } F_{ij}),$$

is nondeg, where  $F_i = \frac{\partial F}{\partial z^i}$ ,  $F_{ij} = \frac{\partial F}{\partial z^i \partial z^j}$  etc. Then

$$\phi : M \rightarrow V, \quad z = (z^1, \dots, z^n) \mapsto (z, F_1, \dots, F_n)$$

is a nondeg. Lagr. immersion and, thus, induces an affine special pseudo-Kähler structure  $(J, g, \nabla)$  on  $M$ .

### Definition

Affine special pseudo-Kähler manifolds as in the above example are called **affine special pseudo-Kähler domains**. The function  $F$  is called a **holomorphic prepotential**.

## Extrinsic construction of special Kähler manifolds III

Since every Lagrangian submanifold of  $(V, \Omega)$  is locally defined by equations  $w_i = F_i(z)$ ,  $i = 1, \dots, n$ , for some hol. function  $F$  and some choice of adapted coordinates  $(z^i, w_i)$ , we obtain:

### Corollary

*Let  $(M, J, g, \nabla)$  be an affine special pseudo-Kähler manifold. Then for every  $p \in M$  there exists a neighborhood  $U$  isomorphic to an affine special pseudo-Kähler domain.*

### Remark

Similar results hold for conical and projective special Kähler manifolds. CASK manifolds are realized as **conical** hol. nondeg. Lagrangian immersions. The corresponding prepotential is defined on a  $\mathbb{C}^*$ -invariant domain  $M \subset \mathbb{C}^n$  and is required to be **homogeneous of degree 2** and to satisfy:  $\sum N_{ij} z^i \bar{z}^j > 0$  and the real symmetric matrix  $(N_{ij})$  has signature  $(1, n - 1)$  on  $M$ .



# Extrinsic construction of special Kähler manifolds IV

Example (complex hyperbolic space as PSK domain)

$$F = \frac{i}{4} \left( (z^0)^2 - \sum_{j=1}^n (z^j)^2 \right)$$

on  $M = \{|z^0|^2 - \sum_{j=1}^n |z^j|^2 > 0\} \subset \mathbb{C}^{n+1}$  is a prepot. for a CASK domain  $(M, J, g, \nabla, \xi)$ . The corresponding PSK domain is  $\mathbb{C}H^n$ .

# Hyper-Kähler manifolds

## Definition

A (pseudo-) hyper-Kähler manifold  $(M, g, J_1, J_2, J_3)$  is a (pseudo-) Riemannian manifold  $(M, g)$  endowed with 3 pairwise anticomm. parallel skew-symm. cx. structures  $J_1, J_2, J_3$  s.t.  $J_3 = J_1 J_2$ .

## Remark

$\implies (M, J_\alpha, g)$  is (pseudo-) Kähler for  $\alpha = 1, 2, 3$ .

## Example

$\mathbb{H}^n = \mathbb{R}^{4n}$  with the Euclidean scalar product  $\langle \cdot, \cdot \rangle$  and  $J_1 = L_i, J_2 = L_j, J_3 = L_k$  is a hyper-Kähler manifold.

# Quaternionic Kähler manifolds I

## Definition

- (i) A **quaternionic structure** on a vector space  $V$  is a subspace  $Q \subset \text{End}(V)$  spanned by three pairwise anticomm. cx. structures  $I, J, K$  s.t.  $K = IJ$ .
- (ii) An **almost quaternionic structure** on a manifold  $M$  is a subbundle  $Q \subset \text{End}(TM)$  such that  $Q_p$  is a quaternionic structure on  $T_pM$  for all  $p$ .  
The bundle  $Q$  is called a **quaternionic structure** if it is parallel for some torsion-free connection.
- (iii) Let  $M$  be a mf. of  $\dim > 4$ . A **quaternionic Kähler structure** on  $M$  is a pair  $(g, Q)$  consisting of a Riem. metric  $g$  and a parallel quaternionic structure  $Q \subset \mathfrak{so}(TM)$ . The triple  $(M, g, Q)$  is called a **quaternionic Kähler (QK) manifold**.

## Remark

If  $\dim M = 4$  in (iii), one has to require in addition  $Q \cdot R = 0$ , which is automatic in higher dimensions.

# Quaternionic Kähler manifolds II

## Fundamental fact

Quaternionic Kähler manifolds are **Einstein**.  $\implies$  3 cases:  
 $Ric = 0$ ,  $Ric > 0$ ,  $Ric < 0$ .

## Relevance to scalar geometry of $N = 2$ theories

**Hyper-Kähler** mfs. = scalar mfs. of **hypermultiplets**,  
**Quaternionic Kähler** mfs. (of  $Ric < 0$ ) = scalar mfs. of  
**supergravity coupled to hypermultiplets** [BW].

## Examples

$Ric = 0$  : Ricci-flat s.c. QK mfs. are **HK**.

$Ric > 0$  : Only known examples of complete QK mfs. of  $Ric > 0$  are the  
**Wolf spaces** = QK symmetric spaces of compact type  
(described below).

- ▶ Simplest example:  $\mathbb{H}P^n$ .

# Quaternionic Kähler manifolds III

## Examples continued

$Ric < 0$  : Known complete QK mfs. of  $Ric < 0$ :

- ▶ QK symm. spaces of noncp. type (dual to Wolf spaces), such as  $\mathbb{H}H^n$ .
- ▶ **Loc. symm. QK mfs.** (including compact examples).
- ▶ **Alekseevsky spaces** (homog. including nonsymm. examples).
- ▶ Deformations of  $\mathbb{H}H^n$ , see [L].
- ▶ New explicit examples obtained using results explained in the next lectures.

## Quaternionic Kähler manifolds IV:

The Wolf spaces can be obtained as follows:

- ▶ Let  $G$  be a cp. s.c. **simple Lie group** and  $\mathfrak{h} \subset \mathfrak{g} = \text{Lie } G$  a Cartan subalgebra,
- ▶  $\mu$  the highest root w.r.t. some system of simple roots and  $\mathfrak{s}_\mu^{\mathbb{C}} = \text{span}\{H_\mu, E_{\pm\mu}\} \subset \mathfrak{g}^{\mathbb{C}}$  the corresponding 3-dim. subalg.
- ▶  $H_\mu \in i\mathfrak{h}$  is normalized such that  $[H_\mu, E_{\pm\mu}] = \pm 2E_{\pm\mu}$ . Then  $\text{ad}_{H_\mu}$  has eigenvalues  $0, \pm 1, \pm 2$  and defines a grading

$$\mathfrak{g}^{\mathbb{C}} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2,$$

where  $\mathfrak{g}_{\pm 2} = \mathbb{C}E_{\pm\mu}$  and  $\mathfrak{g}_0 = \mathbb{C}H_\mu \oplus Z_{\mathfrak{g}^{\mathbb{C}}}(s_\mu^{\mathbb{C}})$ .

- ▶ Put  $\mathfrak{s}_\mu := \mathfrak{g} \cap \mathfrak{s}_\mu^{\mathbb{C}}$ ,  $\mathfrak{k} := \mathfrak{g} \cap \sum_{i=0, \pm 2} \mathfrak{g}_i = \mathfrak{s}_\mu \oplus Z_{\mathfrak{g}}(s_\mu) = N_{\mathfrak{g}}(s_\mu)$ ,

$$\mathfrak{m} := \mathfrak{g} \cap \sum_{i=\pm 1} \mathfrak{g}_i.$$

- ▶ Then  $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$  is a symmetric decomposition, which defines a cp. **QK symm. space**  $M = G/K$ ,  $K = N_G(s_\mu)$ .

# Quaternionic Kähler manifolds V:

## Remarks

- ▶ The invariant quat. structure  $Q$  of the Wolf spaces is defined at  $o = eK \in M = G/K$  as the image of  $s_\mu \cong \mathfrak{sp}(1)$  under the adjoint action on  $\mathfrak{m} \cong T_oM$ .
- ▶ It is skew-symmetric w.r.t. the inv. Riem. metric of  $M$ . The invariance of  $Q$  follows from  $K = N_G(s_\mu)$  and implies that  $Q$  is parallel,
- ▶ since the holonomy group of  $M$  is identified with the isotropy group

$$\text{Hol} = \text{Ad}_K|_{\mathfrak{m}},$$

by the Ambrose-Singer theorem.

- ▶ The noncompact duals  $M^{nc} = G^{nc}/K$  are obtained from the symmetric decomposition  $\mathfrak{g}^{nc} = \mathfrak{k} + i\mathfrak{m}$ .

# Quaternionic Kähler manifolds VI:

The list of Wolf spaces  $M = G/K$

► Classical:

$$\frac{\mathrm{Sp}(n+1)}{\mathrm{Sp}(n)\mathrm{Sp}(1)} = \mathbb{H}P^n = \mathrm{Gr}_1(\mathbb{H}^{n+1}),$$

$$\frac{\mathrm{SU}(n+2)}{\mathrm{S}(\mathrm{U}(n)\mathrm{U}(2))} = \mathrm{Gr}_2(\mathbb{C}^{n+2}),$$

$$\frac{\mathrm{SO}(n+4)}{\mathrm{SO}(n)\mathrm{SO}(4)} = \mathrm{Gr}_4(\mathbb{R}^{n+4}),$$

► Exceptional:

$$\begin{aligned} & \mathrm{G}_2/\mathrm{SO}(4), \quad \mathrm{F}_4/(\mathrm{Sp}(3)\mathrm{Sp}(1)), \\ & \mathrm{E}_6/(\mathrm{SU}(6)\mathrm{Sp}(1)), \quad \mathrm{E}_7/(\mathrm{Spin}(12)\mathrm{Sp}(1)), \quad \mathrm{E}_8/(\mathrm{E}_7\mathrm{Sp}(1)). \end{aligned}$$